

375mm



HUMIC 18

ACTIVE INGREDIENT: Humates and fulvates derived from Leonardite plus 1.8% Potassium

- Improves soil fertility allowing better use of applied nutrients while reducing nutrient leaching
- Improves the ability of soil to exchange nutrients
- Buffers against the negative effects of soil pH and salinity from recycled water
- Improves irrigation water percolation and moisture retention in root zones

CONTENTS
20 L

ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au



DIRECTIONS FOR USE

Shake well before use.

SITUATION	RATE	CRITICAL COMMENTS
Vegetables & other field crops	5-10L/ha	Apply at time of crop flowering, tuber set or heart formation (4-6 wks later) or with soil fertiliser applications.
Perennial crops	5-10L/ha	Apply via drip irrigation just after budburst or first flush or prior to liquid soil applied fertiliser.
Turf establishment Seeding, Sodding, Stolonising	2L/100kg of seed	Seeding: Mix 2L of product and spray on the seeds and mix well. Spread seed out to dry if it appears to be wet.
	20L/ha	Sodding, Stolonising: Apply as a drench with other liquid starter fertilisers at 2L/1000m ² immediately after planting.
Turf maintenance Boom spray or through irrigation system	20L/ha	Apply with normal spraying equipment. Jet size should be bigger than 60 mesh (0.25mm). In order for the product to reach the root zone, the treated area needs to be irrigated after the application. The product can be injected into irrigation systems using a "Dosatron" or similar fertiliser injector equipment.

COMPATIBILITY: OCP HUMIC 18 can be mixed with common plant protection products or fertilisers. Avoid pre-mixing OCP HUMIC 18 with products sensitive to a high pH – tank mixing will not pose any problems. Always add OCP HUMIC 18 at the end into the already prepared plant protection or fertiliser tank mix. Do not mix with calcium-based products. Always conduct a jar compatibility test before mixing with other fertilisers.

PHYTOTOXICITY: OCP HUMIC 18 is non-phytotoxic but has the potential to stain walls and porous surfaces. Wash off after spraying.

SAFETY DIRECTIONS: Avoid contact with eyes. In case of contact with the eyes, rinse with plenty of water. Wash hands and exposed skin with water and soap after work.

STORAGE AND DISPOSAL: OCP HUMIC 18 is completely inert but will absorb moisture over time so keep contents sealed in original container and away from excessively moist environments. Rinse container and dispose of in an environmentally responsible manner.

FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 131 126). If medical advice is needed, have product container or label at hand.

FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 220 770 (ALL HOURS)

SDS: Additional information is listed in the Safety Data Sheet available from the supplier and ocp.com.au.


The OCP drop device is a registered trade mark of Organic Crop Protectants.



BATCH NO:

DOM:

Y20-3517



Job OCP659_Humic 18	Version
Size 375mm x 178mm	Process Flexo Lytho Gravure Offset
Colours C M Y K	Barcode 100% Spot
	FA 17/05/21